

## Yrast and non-yrast spectroscopy of $^{199}\text{Tl}$ using $\alpha$ -induced reactions

Soumik Bhattacharya,<sup>1,2</sup> S. Bhattacharyya,<sup>1,2,\*</sup> R. Banik,<sup>1,2</sup> S. Das Gupta,<sup>3</sup> G. Mukherjee,<sup>1,2</sup> A. Dhal,<sup>1</sup> S. S. Alam,<sup>1,2</sup> Md. A. Asgar,<sup>1,2,†</sup> T. Roy,<sup>1,2</sup> A. Saha,<sup>1,2</sup> S. Nandi,<sup>1,2</sup> T. Bhattacharjee,<sup>1,2</sup> A. Choudhury,<sup>1</sup> Debasish Mondal,<sup>1,2</sup> S. Mukhopadhyay,<sup>1</sup> P. Mukhopadhyay,<sup>1</sup> S. Pal,<sup>1</sup> Deepak Pandit,<sup>1</sup> I. Shaik,<sup>1</sup> and S. R. Banerjee<sup>1</sup>

<sup>1</sup>Variable Energy Cyclotron Centre, I/AF Bidhannagar, Kolkata 700064, India

<sup>2</sup>Homi Bhabha National Institute, Training School Complex, Anushaktinagar, Mumbai-400094, India

<sup>3</sup>Victoria Institution (College), Kolkata 700009, India

(Received 28 March 2018; revised manuscript received 21 August 2018; published 11 October 2018)

The excited states of the  $^{199}\text{Tl}$  nucleus have been studied by using the light ion induced fusion evaporation reaction  $^{197}\text{Au}(\alpha, 2n)^{199}\text{Tl}$  at 30 MeV of beam energy by  $\gamma$ -ray spectroscopic methods. VECC Array for NUclear Spectroscopy (VENUS) has been used to detect the prompt  $\gamma$  rays. Level scheme of  $^{199}\text{Tl}$  has been significantly improved and extended with the placement of 53 new transitions. The yrast  $\pi h_{9/2}$  band has been extended in this nucleus beyond the band crossing. Several new near- and non-yrast band structures have also been identified. It has been observed that in case of the three-quasiparticle structures, a different configuration involving negative parity neutron orbitals becomes yrast for heavier Tl isotopes with  $N \geq 118$  in contrast to the involvement of the neutron  $i_{13/2}$  orbital for lighter Tl isotopes. It was possible to identify both the yrast and the non-yrast states corresponding to these configurations in the present work. The observed band structures have been interpreted in light of the systematics of the neighboring odd mass Thallium nuclei. Total Routhian surface calculations have been performed to study the deformation and shape changes as a function of rotational frequency in this nucleus.

DOI: [10.1103/PhysRevC.98.044311](https://doi.org/10.1103/PhysRevC.98.044311)

### I. INTRODUCTION

Thallium isotopes, with only one proton below the  $Z = 82$  shell closure, are the most appropriate candidates to extend our knowledge on the polarizing effect of the high- $j$  proton intruder orbitals (e.g.,  $\pi h_{9/2}, \pi i_{13/2}$ ) or high- $j$  neutron orbitals (e.g.,  $\nu i_{13/2}$ ) on the shape of a nucleus, which are otherwise near-spherical at their ground state. Several interesting and exotic phenomena, like magnetic rotational bands [1] and chiral bands [2–4] have been reported in the thallium isotopes around the  $A \sim 200$  mass region. It is, therefore, interesting to investigate the interplay of the single particle structures (involving high- $j$  orbitals) and the collectiveness of the underlying core that generates the above exotic phenomena, below the  $Z = 82$  shell closure region [5–7]. It is known that the high- $j$  orbitals, specially,  $\pi h_{9/2}$  and  $\nu i_{13/2}$ , induce oblate shape in the nuclei in this region [8]. In the case of the odd- $A$  Tl nuclei, a wide diversity of shapes and structures have been observed from superdeformed structures in neutron deficient  $^{191,193,195}\text{Tl}$  [9–11] to weakly deformed oblate band structures in  $^{191,193,195,197,201}\text{Tl}$  [12–15], depending on the neutron Fermi level and the excitation energy. In the heavier isotope  $^{205}\text{Tl}$ , which is close to the doubly magic  $^{208}\text{Pb}$  nucleus, excited states corresponding to the octupole core excitation was also observed [16]. Although deformed shapes based on the  $\pi i_{13/2}$  intruder orbital have been observed for the lighter Tl isotopes,

no such band structure has been reported in heavier odd-mass thallium isotopes. A survey of the excitation energies of these states in the heavier isotopes indicates that it becomes more and more non-yrast with the increase in neutron number [17]. In the neighboring isotope  $^{197}\text{Tl}$ , an excited state corresponding to the  $\pi i_{13/2}$  orbital has been reported but no band structure was observed on top of this state [18].

It is interesting to note that the shape polarizing effect of the  $\pi h_{9/2}$  orbital continues to generate deformed band structures for the isotopes  $^{200,201}\text{Tl}$ , as reported in our previous work [15,19]. However, the three-quasiparticle configuration, observed after the band crossing of the  $\pi h_{9/2}$  oblate band, in the  $N = 120$  isotope  $^{201}\text{Tl}$  was observed to be different with smaller gain in aligned angular momentum compared to the other lighter isotopes. This is possibly because of the fact that the neutron Fermi level moves up and away from the  $\nu i_{13/2}$  orbital with the increase in neutron number. For spherical shape, the neutron Fermi level is expected to be situated around the  $3p_{3/2}$  orbital above the  $i_{13/2}$  orbital for  $N \geq 114$ . However, for oblate deformation, as it is the case for the Tl isotopes, it would move up to lie close to the  $2f_{5/2}$  orbital for  $N \geq 118$ . Therefore, it is important to study the band crossing behavior of  $^{199}\text{Tl}$  to understand the relative position of the  $\nu i_{13/2}$  orbital.

The available information on the excited states in  $^{199}\text{Tl}$  is very scarce and limited to a few low-lying states which precludes one to get any idea about the  $\nu i_{13/2}$  alignment. One of the first measurements of the excited states in  $^{199}\text{Tl}$  was performed way back in 1970 by Newton *et al.* [20], with a few Ge(Li) detectors. Although only a few states could be identified in that work, but importantly, the deformed nature

\*Corresponding author: sarmi@vecc.gov.in

†Present address: Prabhat Kumar College, Contai, Purba Medinipur, West Bengal, India.

- [13] J. O. Newton, F. S. Stephens, and R. M. Diamond, *Nucl. Phys. A* **236**, 225 (1974).
- [14] R. M. Lieder *et al.*, *Nucl. Phys. A* **299**, 255 (1978).
- [15] S. DasGupta, S. Bhattacharyya, H. Pai, G. Mukherjee, S. Bhattacharya, R. Palit, A. Shrivastava, A. Chatterjee, S. Chanda, V. Nanal, S. K. Pandit, S. Saha, J. Sethi, and S. Thakur, *Phys. Rev. C* **88**, 044328 (2013).
- [16] J. Wrzesiński *et al.*, *Eur. Phys. J. A* **20**, 57 (2003).
- [17] R. B. Firestone *et al.*, *Table of Isotopes* (John Wiley and Sons, New York, 1999).
- [18] H. Pai, G. Mukherjee, S. Bhattacharya, C. Bhattacharya, S. Bhattacharyya, T. Bhattacharjee, S. Chanda, S. Rajbanshi, A. Goswami, M. R. Gohil, S. Kundu, T. K. Ghosh, K. Banerjee, T. K. Rana, R. Pandey, G. K. Prajapati, S. R. Banerjee, S. Mukhopadhyay, D. Pandit, S. Pal, J. K. Meena, P. Mukhopadhyay, and A. Choudhury, *Phys. Rev. C* **88**, 064302 (2013).
- [19] S. Bhattacharya *et al.*, *Phys. Rev. C* **95**, 014301 (2017).
- [20] J. O. Newton *et al.*, *Nucl. Phys. A* **148**, 593 (1970).
- [21] R. E. Doebler and Wm. C. McHarris, *Phys. Rev. C* **2**, 2422 (1970).
- [22] N. Marginean *et al.*, *Eur. Phys. J. A* **46**, 329 (2010).
- [23] S. Bhattacharya *et al.*, DAE-BRNS Symp. Nucl. Phys. **61**, 98 (2016).
- [24] <http://www.tifr.res.in/~pell/lamps.html>
- [25] R. K. Bhowmik, S. Muralithar, and R. P. Singh, Proc. DAE Symp. Nucl. Phys. **44B**, 422 (2001).
- [26] D. C. Radford, *Nucl. Instrum. Methods Phys. Res. A* **361**, 297 (1995).
- [27] A. Kramer-Flecken *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **275**, 333 (1989).
- [28] K. Starosta *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **423**, 16 (1999).
- [29] Ch. Droste *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **378**, 518 (1996).
- [30] B. Singh, *Nucl. Data Sheets* **108**, 79 (2007).
- [31] R. M. Diamond and F. S. Stephens, *Nucl. Phys. A* **45**, 632 (1963).
- [32] C. Günther *et al.*, *Phys. Rev. C* **15**, 1298 (1977).
- [33] K. H. Hicks, T. E. Ward, J. Wiggins, C. A. Fields, and F. W. N. deBoer, *Phys. Rev. C* **25**, 2710 (1982).
- [34] L. L. Collins *et al.*, Diss. Abst. Int. **39B**, 3834 (1979).
- [35] M. G. Slocombe, J. O. Newton, and G. D. Dracoulis, *Nucl. Phys. A* **275**, 168 (1977).
- [36] D. Royer *et al.*, *Nucl. Phys. A* **158**, 516 (1970).
- [37] J. A. Becker, R. G. Lanier, L. G. Mann, G. L. Struble, K. H. Maier, L. E. Ussery, W. Stoffl, T. W. Nail, R. K. Sheline, J. A. Cizewski, B. H. Erkkila, and J. Blomqvist, *Phys. Rev. C* **29**, 1268 (1984).
- [38] H. Beuscher, P. Jahn, R. M. Lieder, and C. Mayer-Boricke, *Z. Phys. C* **247**, 383 (1971).
- [39] Z. Plajner, J. Frana, I. Rezanka, A. Spalek, and M. Fiser, *Z. Phys. C* **233**, 122 (1971).
- [40] K. Farzine and H. V. Buttlar, *Z. Phys. C* **270**, 155 (1974).
- [41] A. J. Haverfield, H. T. Easterday, and J. M. Hollander, *Nucl. Phys. A* **64**, 379 (1965).
- [42] W. Nazarewicz *et al.*, *Nucl. Phys. A* **435**, 397 (1985).
- [43] W. Nazarewicz *et al.*, *Nucl. Phys. A* **512**, 61 (1990).
- [44] J. Dudek, Z. Szymanski, and T. Werner, *Phys. Rev. C* **23**, 920 (1981).
- [45] G. Mukherjee *et al.*, *Nucl. Phys. A* **829**, 137 (2009).

# Spectroscopy of a tetrahedral doubly magic candidate nucleus $^{160}_{70}\text{Yb}_{90}$

A Saha<sup>1,2,12</sup>, T Bhattacharjee<sup>1,2</sup>, D Curien<sup>3</sup>, J Dudek<sup>3,4</sup>, I Dedes<sup>4</sup>, K Mazurek<sup>5</sup>, A Góźdż<sup>4</sup>, S Tagami<sup>6</sup>, Y R Shimizu<sup>6</sup>, S R Banerjee<sup>1,2</sup>, S Rajbanshi<sup>7,13</sup>, A Bisoi<sup>7,14</sup>, G De Angelis<sup>8</sup>, Soumik Bhattacharya<sup>1,2</sup>, S Bhattacharyya<sup>1,2</sup>, S Biswas<sup>9</sup>, A Chakraborty<sup>10,15</sup>, S Das Gupta<sup>1,16</sup>, B Dey<sup>1,2,17</sup>, A Goswami<sup>7</sup>, D Mondal<sup>1,2</sup>, D Pandit<sup>1,2</sup>, R Palit<sup>9</sup>, T Roy<sup>1,2</sup>, R P Singh<sup>11</sup>, M Saha Sarkar<sup>7</sup>, S Saha<sup>9</sup> and J Sethi<sup>9</sup>

<sup>1</sup> Variable Energy Cyclotron Centre, IN-700064 Kolkata, India

<sup>2</sup> Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, IN-400094 Mumbai, India

<sup>3</sup> Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

<sup>4</sup> Institute of Physics, Marie Curie-Skłodowska University, PL-20031 Lublin, Poland

<sup>5</sup> The Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, 2 ul. Radzikowskiego 152, PL-31342 Kraków, Poland

<sup>6</sup> Department of Physics, Faculty of Sciences, Kyushu University, Fukuoka JP-8190359, Japan

<sup>7</sup> Saha Institute of Nuclear Physics, IN-700064 Kolkata, India

<sup>8</sup> INFN, Laboratori Nazionali di Legnaro, IT-35020 Legnaro, Italy

<sup>9</sup> Tata Institute of Fundamental Research, IN-400005 Mumbai, India

<sup>10</sup> Department of Physics, Krishnath College, IN-742101 West Bengal, India

<sup>11</sup> Inter University Accelerator Centre, IN-110067 New Delhi, India

E-mail: [btumpa@vecc.gov.in](mailto:btumpa@vecc.gov.in)

Received 9 June 2018, revised 31 January 2019

Accepted for publication 8 February 2019

Published 8 March 2019



CrossMark

## Abstract

The decay scheme of  $^{160}\text{Yb}$  nucleus populated in the  $^{148}\text{Sm}(^{16}\text{O}, 4\text{n})^{160}\text{Yb}$  reaction at 90 MeV has been studied. The gamma-coincidence data have been collected by using Indian National Gamma Array (INGA) composed of twenty

<sup>12</sup> Present Address: Physics Department, Indian Institute of Technology Bombay, Powai, IN-400 076 Mumbai, India.

<sup>13</sup> Present Address: Dum Dum Motijheel College; 1, Motijheel Avenue; Dum Dum; IN-700074 Kolkata, India.

<sup>14</sup> Present Address: Indian Institute of Engineering Science and Technology, Shibpur, IN-711103 West Bengal, India.

<sup>15</sup> Present Address: Siksha Bhavana, Visva-Bharati, Santiniketan 731 235, West Bengal, India.

<sup>16</sup> Present Address: Victoria Institution (College), 78 B, A P C Road, IN-700 009 Kolkata, India.

<sup>17</sup> Present Address: Saha Institute of Nuclear Physics, IN-700064 Kolkata, India.

- [38] Droste C, Rohoziński S G, Starosta K, Morek T, Srebrny J and Magierski P 1996 *Nucl. Instrum. Methods A* **378** 518
- [39] Starosta K *et al* 1999 *Nucl. Instrum. Methods A* **423** 16
- [40] Jones P M, Wei L, Beck F A, Butler P A, Byrski T, Duchêne G, de France G, Hannachi F, Jones G D and Kharraja B 1995 *Nucl. Instrum. Methods A* **362** 556
- [41] Bhattacharjee T *et al* 2008 *Phys. Rev. C* **78** 024304
- [42] Reich C W 2005 *Nucl. Data Sheets* **105** 557
- [43] Experimental Unevaluated Nuclear Data List Search and Retrieval (XUNDL) <http://nndc.bnl.gov/ensdf/ensdf/xndl.jsp>, (Last updated: 02 February 2018)
- [44] Gavron A 1980 *Phys. Rev. C* **21** 230
- [45] Campbel D B 2004 Angular Momentum Induced Shape Changes in the Rare-Earth Nuclei 152,153Gd and 159,160Yb *PhD Thesis* unpublished <http://diginole.lib.fsu.edu/islandora/object/fsu%3A185022> Florida State University
- [46] Saha A *et al* 2018 *Phys. Scr.* **93** 034001
- [47] Reviol W *et al* 2006 *Phys. Rev. C* **74** 044305
- [48] Zimba G L *et al* 2016 *Phys. Rev. C* **94** 054303
- [49] Dudek J, Dedes I, Yang J, Baran A, Curien D, Dickel T, Góźdż A, Rouvel D and Wang H-L 2019 *Acta Phys. Pol.* in print **50**



shinjinee dasgupta &lt;shinjinee14@gmail.com&gt;

## Schedule of INGA Experiment at VECC

2 messages

Chandana Bhattacharya &lt;chandana@vecc.gov.in&gt;

Fri, Oct 26, 2018 at 6:39 PM

To: r.palit1@gmail.com, somsundarm@gmail.com, ritwika.chakrabarti@gmail.com, sujit.tandel@cbs.ac.in, shinjinee14@gmail.com

Cc: Amitava Roy &lt;amitav@vecc.gov.in&gt;, Prodyut Sankar Chakraborty &lt;prodyut@vecc.gov.in&gt;, Tapas Bandyopadhyay &lt;tapas@vecc.gov.in&gt;, asimananda.goswami@saha.ac.in, Sandeep Ghugre &lt;ssg.iuc@gmail.com&gt;, Sarmishtha Bhattacharya &lt;sarmi@vecc.gov.in&gt;, Gopal Mukherjee &lt;gopal@vecc.gov.in&gt;, Soumik Bhattacharya &lt;oumik@vecc.gov.in&gt;

Dear All,

Please find attached herewith the schedule of your experiment using INGA- facility at VECC. Please note that the beam time allotted includes radioactivity/ decay run, calibration/efficiency run.

You are requested to send an acknowledgement at the earliest.

Please also note that, for smooth running of the experiment, a local experimental coordinator (LEC) has been assigned for each experiment.

You are requested to send certain information as per attached documents to the LEC, e-mails given below.

LEC:

Dr. Gopal Mukherjee: [gopal@vecc.gov.in](mailto:gopal@vecc.gov.in)Dr. Sarmishtha Bhattacharyya: [sarmi@vecc.gov.in](mailto:sarmi@vecc.gov.in)Shri Soumik Bhattacharya: [soumik@vecc.gov.in](mailto:soumik@vecc.gov.in)

With best regards

Chandana bhattacharya

Dr.(Mrs.) Chandana Bhattacharya  
 Head, Experimental Nuclear Physics Division  
 Physics Group  
 Variable Energy Cyclotron Centre  
 1/AF, Bidhan Nagar, Kolkata  
 India  
 Ph:+91-33-2318-2303  
 Fax:+91-33-2334-6871

---

### 3 attachments

[radiation worker Visitor form \(2\).pdf](#)  
21K

[Info\\_user.pdf](#)  
10K

[Beamtime\\_Schedule\\_8\\_11\\_18-7\\_12\\_18.pdf](#)  
58K

---

shinjinee dasgupta <shinjinee14@gmail.com>  
 To: chandana@vecc.gov.in

Sat, Oct 27, 2018 at 3:16 PM

Respected Madam,

10/6/23, 10:11 PM

Gmail - Schedule of INGA Experiment at VECC

Thank you for scheduling the experiment and I want to confirm my participation.

With regards,  
Shinjinee

[Quoted text hidden]