
Fwd: CATENA21956R2 - Editor's decision: accepted

Rajkumar Bhattacharya <rajgeovu10@gmail.com>

Thu, Jun 13, 2024 at 6:20 PM

To: "Nilanjana Das Faculty, Geography" <nilanjana_vu@mail.vidyasagar.ac.in>

----- Forwarded message -----

From: **Catena** <em@editorialmanager.com>

Date: Thu, 13 Jun 2024, 18:10

Subject: CATENA21956R2 - Editor's decision: accepted

To: **Raj Kumar Bhattacharya** <rajgeovu10@gmail.com>

Dear Dr. Bhattacharya,

I am pleased to inform you that the manuscript "Linkages between gully erosion susceptibility and hydrological connectivity in Tropical sub-humid river basin: Application of Machine learning algorithm and Connectivity Index model" (Dr. Raj Kumar Bhattacharya) has now been accepted by the editor for publication.

Your accepted manuscript will now be transferred to our production department and work will begin on creation of the proof. If we need any additional information to create the proof, we will let you know. If not, you will be contacted again in the next few days with a request to approve the proof and to complete a number of online forms that are required for publication.

We appreciate and value your contribution to Catena. We regularly invite authors of recently published manuscript to participate in the peer review process. If you were not already part of the journal's reviewer pool, you have now been added to it. We look forward to your continued participation in our journal, and we hope you will consider us again for future submissions.

We encourage authors of original research papers to share the research objects – including raw data, methods, protocols, software, hardware and other outputs – associated with their paper. More information on how our open access Research Elements journals can help you do this is available at https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec_em_research_elements_email.

Kind regards,


Zhongping Lai
Editor
Catena

Comments from the Editors and Reviewers:

At Elsevier, we want to help all our authors to stay safe when publishing. Please be aware of fraudulent messages requesting money in return for the publication of your paper. If you are publishing open access with Elsevier, bear in mind that we will never request payment before the paper has been accepted. We have prepared some guidelines (<https://www.elsevier.com/connect/authors-update/seven-top-tips-on-stopping-apc-scams>) that you may find helpful, including a short video on Identifying fake acceptance letters (<https://www.youtube.com/watch?v=o5l8thD9XtE>). Please remember that you can contact Elsevier's Researcher Support team (<https://service.elsevier.com/app/home/supporthub/publishing/>) at any time if you have questions about your manuscript, and you can log into Editorial Manager to check the status of your manuscript (https://service.elsevier.com/app/answers/detail/a_id/29155/c/10530/supporthub/publishing/kw/status/).

#AU_CATENA#

To ensure this email reaches the intended recipient, please do not delete the above code



Linkages between gully erosion susceptibility and hydrological connectivity in Tropical sub-humid river basin: Application of Machine learning algorithms and Connectivity Index

Raj Kumar Bhattacharya^{a,1,*}, Nilanjana Das Chatterjee^{b,2}, Kousik Das^{b,3}

^a Department of Geography, Victoria Institution (College), 78B, Acharya Prafulla Chandra Road, Kolkata, 700009, India

^b Department of Geography, Vidyasagar University, Midnapore, 721102, West Bengal, India

ARTICLE INFO

Keywords:

Land degradation
Sediment detachment
Gully susceptibility
Hydrological connectivity
Sustainable land uses policy

ABSTRACT

Hydrological connectivity from upslope to downslope of valley floor and main channel, triggered the gully initiation and associated land degradation continue occurring off-site erosion as considered most effective drivers on potential sediment detachment. Present study attempted to identify the linkage between gullies erosion susceptibility (GES) and hydrological connectivity pathway in sub-tropical humid river basin Kangsabati (KRB) using four machine-learning algorithms (MLALs) such as Random Forest (RF), Support Vector Machine (SVM), Extreme Gradient Boosting (XGB), Artificial Neural Network (ANN) for GES mapping, and connectivity index (IC) for hydrological connectivity mapping. Thirty-five controlling factors were selected using Boruta's approach to produce GES mapping, while frequency ratio (FR) was applied to determine the significant role in each individual class of controlling factors on degree of gully susceptibility. To achieve the efficiency of using MLALs, AUC of ROC including sensitivity, specificity, accuracy, F, and Kappa index were employed to compare in each model. In testing datasets, AUC values reveals that RF (0.99) and XGB (0.99) were well performed and predicted to GES followed by ANN (0.97) and SVM (0.87). FR depicts the most contributing factors of barren land and laterite followed by rainfall erosivity, degraded forest, single crop, and elevation to GES. IC result showed that values range (-11.52 to 0.49) address the three connectivity categories i.e. not connected (NC), gully connected but not reach (CNR), and gully well connected (WC). Correlation analysis clarified that double crop ($R = 0.82$), topographical wetted index ($R = 0.77$) and slope ($R = 0.044$) are prolonged WC for large downslope in north-western sub-basins of upper catchment and western, eastern sub-basins of lower catchment, while dense forest ($R = -0.71$) and vegetation cover ($R = -0.82$) forms NC gullies and CNR to channel due to interrupted upslope in central sub-basins of upper and lower catchment. Finally, research findings could provide to take strategy for sustainable land uses policy.

1. Introduction

Land is an essential component of the earth's surface as provided survival opportunities and fulfils the necessary demand of human beings through the various way likes agriculture, forest, industry, residence, etc. (Saha et al., 2021); however, man is regarded as a crucial responsible agent for modification, management and intended way of land resource utilization (Turner et al., 1995). In contrary, sustainable land use arrangements always preserve the basic fundamental functional

processes amongst the biosphere community, increase the quality of life and maintain their positive interaction with the environment (Yao et al., 2018). In India, disorganized land use arrangement in the entire Chotanagpur plateau of western West Bengal has been facing under sub-aerial denudation process caused by intensive water induced soil erosion through the gully formation and its development (Saha et al., 2021).

Generally, gullies are rapidly developed through the shifting of underlying unconsolidated material from uplands to valley bottoms with

* Corresponding author.

E-mail addresses: rajgeovu0@gmail.com (R. Kumar Bhattacharya), nilanjana_vu@mail.vidyasagar.ac.in (N. Das Chatterjee).

¹ ORCID- <https://orcid.org/0000-0002-1102-0088>.

² ORCID- <https://orcid.org/0000-0001-9436-2173>.

³ ORCID- <https://orcid.org/0000-0001-9948-1577>.